

Primary Author: Alessandra Diioia, PharmD; Melanie Perry, PharmD
Previous Authors: Kathleen Relihan, MD, Amanda Giancarelli, PharmD, CNSC, Brandon Hobbs, PharmD
Editor: Michael L. Cheatham, MD
Approved: 1/7/03, 10/17/06, 11/1/11, 7/29/15, 1/26/19, 4/8/21 Last revision date: 12/27/2022

SUMMARY

The splenectomized patient should be vaccinated to decrease the risk of overwhelming post-splenectomy sepsis (OPSS) by organisms such as *Streptococcus pneumoniae*, *Haemophilus influenzae* type B, and *Neisseria meningitidis*. Patients should be educated prior to discharge on the risk of OPSS, their immunocompromised state and the need for booster vaccinations. The need for prompt medical attention if fevers or concerning symptoms develop should be instilled in patients to reduce the morbidity and mortality of post-splenectomy infection.

RECOMMENDATIONS

- **Level 1**
 - None
- **Level 2**
 - Non-elective splenectomy patients should be vaccinated 14 days post-splenectomy or at the time of hospital discharge, whichever comes first.
 - Asplenic patients should be revaccinated at the appropriate time interval for each vaccine (Table 1).
 - Following splenic embolization, splenic immune function is retained, and routine vaccination is not indicated.
- **Level 3**
 - Elective splenectomy patients should be vaccinated at least 14 days prior to operation.
 - Asplenic or immunocompromised patients (with an intact but nonfunctional spleen) should be vaccinated as soon as the diagnosis is made.
 - When adult vaccination is indicated, the following FIVE vaccinations should be administered: *
 - Pneumococcal vaccine naïve: 20-valent pneumococcal conjugate vaccine (PCV20) alone or 15-valent pneumococcal conjugate vaccine (PCV15) followed by 23-valent pneumococcal (PPSV23) ≥ 8 weeks later
 - Previous PPSV23 vaccination: PCV20 alone or PCV15 ≥ 1 year after PPSV23
 - Previous PCV13 vaccination: benefits of providing PCV20 or PCV15 have not been evaluated, adults should complete previously recommended PPSV23 series. If the PPSV23 series has not been completed, one dose of PCV20 may be used if PPSV23 is not available
 - MenACWY (Menveo®), two doses, given at least two months apart
 - MenB-FHbp (Trumenba®; three-dose series) at 0, 2, and 6 months OR MenB-04C (Bexsero®; two dose series) at least one month apart
 - *Haemophilus influenzae* b vaccine (Hiberix®)
 - Influenza vaccine annually
 - Pediatric vaccination should be performed according to the recommended pediatric dosage and vaccine types with special consideration made for children less than 2 years of age

*A patient information sheet of post-splenectomy vaccines to provide to patients is available on the last page of this guideline.

LEVEL OF RECOMMENDATION DEFINITIONS

- **Level 1:** Convincingly justifiable based on available scientific information alone. Usually based on Class I data or strong Class II evidence if randomized testing is inappropriate. Conversely, low quality or contradictory Class I data may be insufficient to support a Level I recommendation.
- **Level 2:** Reasonably justifiable based on available scientific evidence and strongly supported by expert opinion. Usually supported by Class II data or a preponderance of Class III evidence.
- **Level 3:** Supported by available data, but scientific evidence is lacking. Generally supported by Class III data. Useful for educational purposes and in guiding future clinical research.

DISCLAIMER: These guidelines were prepared by the Department of Surgical Education, Orlando Regional Medical Center. They are intended to serve as a general statement regarding appropriate patient care practices based on the medical literature and clinical expertise at the time of development. They should not be considered to be accepted protocol or policy, nor are intended to replace clinical judgment or dictate care of individual patients.

INTRODUCTION

Blunt abdominal trauma commonly injures the spleen resulting in either irreparable parenchymal disruption (necessitating removal of the injured organ) or devascularization of varying degrees. Non-operative management may avoid splenectomy, but can also result in functional asplenia if the devascularization is extensive. Splenic injury can also be managed by splenic embolization. Elective splenectomy may be indicated for specific primary disease of the spleen or symptomatic splenomegaly.

Loss of functional splenic tissue places such individuals at high risk for infection by encapsulated organisms such as *Haemophilus influenzae* type b, *Neisseria meningitidis* and *Streptococcus pneumoniae*. Although the risk of fulminant septicemia or meningitis due to such organisms appears to be less in the adult population (by virtue of prior exposure to these bacteria), overwhelming post-splenectomy sepsis (OPSS) remains a significant concern in the asplenic patient (1).

The incidence of OPSS is estimated to occur in 0.05% to 2% of splenectomized patients (2). It may develop immediately or as late as 65 years post-splenectomy (2-4). Most OPSS occurs during the first two years following splenectomy. Mortality associated with OPSS is significant and reported to be as high as 50% (4,5). OPSS incidence reduction is dependent upon education of both the patient and physician about the risks, prevention, and rapid recognition of the asplenic individual when infection may be present (2,5-7).

Reduced post-splenectomy levels of opsonins, splenic tuftsin, and immunoglobulin M (IgM) (which promote phagocytosis of particulate matter and bacteria), hinder the body's ability to clear encapsulated organisms (4,6,7). Vaccination, to impart immunity against such infections, is commonly performed despite the absence of Class I or Class II data to support its efficacy. As 50 to 90% of OPSS infections are secondary to *Streptococcus pneumoniae* infection, the polyvalent pneumococcal vaccine has been the most administered post-splenectomy vaccine. In recent years, the meningococcal and *Haemophilus influenzae* type b vaccines have also been advocated (2-17).

Timing of vaccine administration following splenectomy has been a topic of longstanding debate. Two major concerns include the patients' immunogenicity during the perioperative period and the impaired immune function of critically ill patients (2,13,16,17). The patient's present state of health should be considered prior to administration of post-splenectomy vaccines. In patients with moderate to severe acute illness, vaccination should be delayed until the illness has resolved. This minimizes adverse effects of the vaccine which could be more severe in the presence of illness or could confuse the patient's clinical picture (such as a post-vaccine fever) (16).

With increased use of splenic embolization to manage splenic injuries nonoperatively, there has been growing interest in which patients need vaccinations and which have enough preserved splenic parenchyma to not require vaccination. The Advisory Committee on Immunization Practices (ACIP) recommends that asplenic patients with 50% or more of the splenic mass lost should be viewed as if they are asplenic and should receive vaccinations (18). However, there is no single test available for measuring residual splenic function to determine if vaccinations are indicated to prevent OPSS infections. More recent evidence suggests that splenic function is preserved after embolization and routine post-splenectomy vaccination may not be necessary (19-25). Additionally in August 2022, the Eastern Association for the Surgery of Trauma (EAST) provided recommendations against routine post-splenectomy vaccinations in adult trauma patients who have undergone angioembolization due to retained immune function (26).

Prior to 2021, the only two pneumococcal vaccines approved for use in adult patients were PCV13 and PPSV23. In 2021, the PCV20 and PCV15 were licensed by the Food and Drug Administration (FDA) for adults aged ≥ 18 years (27). On October 20, 2021, the ACIP recommended use of PCV20 alone or PCV15 in series with PPSV23 for all adults ≥ 65 years, and for adults ages 19-64 years with certain underlying medical conditions or other risk factors who have not previously received a PCV or whose previous vaccination history is unknown (27). In 2022, the FDA expanded the use of PCV15 to infants and children aged 6 weeks and older (28). PCV20 helps protect against the 20 *S. pneumoniae* serotypes in the vaccine, including 7 additional serotypes compared to PCV13 (29). PCV15 contains pneumococcal polysaccharide serotypes 22F and 33F in addition to the PCV13 serotypes (27). PPSV23 is a polyvalent vaccine of purified capsular polysaccharides from 23 strains of *Streptococcus pneumoniae* (30). Based on clinical trials, the immunogenicity and safety of PCV20 alone or PCV15 in series with PPSV23 were comparable to PCV13 alone or in series with PPSV23. Additionally, by simplifying pneumococcal vaccine recommendations, cost-effectiveness studies demonstrated cost-saving for all adults ≥ 65 years old (27).

In light of two serogroup B meningococcal (MenB) disease outbreaks on college campuses in 2013, the FDA granted Breakthrough Therapy designations to two MenB vaccines, MenB-FHbp (Trumenba®) and MenB-4C (Bexsero®). Both were approved for ages 10-25 years and are recommended by the ACIP for prevention of serogroup B meningococcal disease in all asplenic patients ≥10 years (31). The MenB-FHbp vaccine is composed of two recombinant factor H binding proteins from *N. meningitidis* serogroup B. It is administered as a three-dose series at 0, 2, and 6 months (32). MenB-4C is composed of three recombinant proteins and outer membrane vesicles. It is administered as a two-dose series given at least one month apart (24). Due to the low incidence of serogroup B meningococcal disease, the efficacy of both vaccines was evaluated in immunogenicity studies and determined based on the proportion of patients who achieved ≥4-fold increase in complement activity against serogroup B strains (31).

All of the vaccines can cause adverse reactions which are generally self-limiting and resolve 24-72 hours after administration. Pneumococcal vaccines may cause a transient and self-limited fever in vaccinated patients, as well as pain and redness at the site for 1-2 days. A hypersensitivity reaction can occur at the injection site of the *Haemophilus influenzae* type b vaccine along with occasional fever, aches, and malaise. The meningococcal vaccines can all cause headaches, fatigue, malaise and injection site reactions.

Another important aspect to consider when managing patients post splenectomy or with evidence of asplenia is long term follow up and need for revaccination. Several studies have shown that patients have poor information recall on inpatient vaccination and low revaccination rates (33). At the end of this guideline is a sample form for patients and clinicians to monitor vaccination and revaccination status.

Table 1: Available post-splenectomy vaccines

Vaccine	Dose	Route	Revaccination
20-valent pneumococcal conjugate vaccine (PCV20, Prevnar 20™)	0.5 mL	IM	None
15-valent pneumococcal (PCV15, Vaxneuvance™)	0.5 mL	IM	None
23-valent pneumococcal (PPSV23, Pneumovax® 23)	0.5 mL	IM or SC	None
Meningococcal/diphtheria conjugate (MenACWY, Menveo® or MenQuadfi®)	0.5 mL	IM	At 2 months & every 5 yrs
Serogroup B meningococcal (MenB-FHbp, Trumenba™)	0.5 mL	IM	At 2 and 6 mo
Serogroup B meningococcal (MenB-4C, Bexsero®)	0.5 mL	IM	Once at ≥1 mo
<i>Haemophilus influenzae</i> b conjugate (Hiberix®)	0.5 mL	IM	None
Influenza	0.5 mL	IM	One dose annually

LITERATURE REVIEW

Two Class I studies have demonstrated that the polyvalent pneumococcal vaccine results in the highest antibody titers, for the most common serotypes, when administered at least 14 days post-splenectomy (16,17). These prospective, randomized trials evaluated the efficacy of the vaccine when administered at 1-, 7-, 14-, and 28-days post-splenectomy. As these trials were designed to demonstrate the immunogenicity of the vaccines and not the prevention of OPSS, they can only be used to advocate *timing* of vaccination.

In 2004, Landgren et al. published a prospective study on antibody response to repeated vaccination. This study included 28 (out of 311) post-trauma splenectomized patients. Their results showed that time between splenectomy and first pneumococcal vaccine was not associated with pre-vaccination, peak or follow-up antibody levels. Twenty-five of the 28 trauma patients received their first vaccine post-splenectomy. A major limitation of this study is that the time from splenectomy to first vaccine was only documented in 24% of the cases, yet they claimed that timing had no effect (34-35). Similarly, Grimfors et al. conducted a longitudinal study of 173 patients (33 trauma) for three years. Pneumococcal antibody responses declined to pre-treatment values at three years in all groups. They also found no correlation between the interval from splenectomy to vaccination and response to vaccination. The data to support this conclusion was not published (36).

Schreiber et al. published a study in rats in 1998 looking at timing of vaccination and subsequent ability to survive pneumococcal challenge. There was no difference between the groups of rats if they were vaccinated on post-operative day 1, 7, or 42 and their ability to survive a pneumococcal challenge (37). In another study, Werner et al.

looked at the effect of perioperative hypovolemic shock and response to vaccination and found no difference if the splenectomized rats were vaccinated on post-operative day 1, 7, or #28. Both studies raise the question of whether delaying vaccination for fourteen days as suggested in the Shatz et al. studies is necessary (38). Further human studies are needed to address timing of post-splenectomy vaccines.

Class II data supports the vaccination of asplenic patients based on studies of the spleen's role in immune function and its ability to provide defense against encapsulated organisms (5). Current Center for Disease Control (CDC) recommendations for post-splenectomy vaccinations include PCV20 alone or PCV15 in series with PPSV23, the meningococcal MenACWY and either MenB-FHbp or MenB-4C, and the *Haemophilus influenzae* type b vaccine (12-14,27,35-46). All four of the initial vaccines may be administered simultaneously (15,31).

The ACIP suggests patients with <50% splenic function should be vaccinated to prevent OPSS. However, how to manage patients who undergo splenic arterial embolization (SAE) is not well known. For SAE, patients can have a partial or total embolization depending upon the degree of injury. Often, clinicians vaccinate those who had total SAE only. A systematic review from the Netherlands by Schimmer et al. found that 11 out of 12 studies evaluating the long-term effects of SAE found that there was preserved splenic function (47). Various studies have evaluated splenic function after SAE, concluding that immunocompetence of the spleen is preserved (20-23). However, there is no one specific test available which can demonstrate if splenic function is preserved, and the majority of studies performed are retrospective with a small sample size. A prospective study by Olthof et al. comparing splenic function of embolized patients to both healthy controls and those who underwent splenectomy, found that the median vaccine specific antibody response of SAE patients did not differ significantly from healthy controls (24). Foley et al. published a prospective study examining the differences in IgM Memory B cells after splenectomy, distal SAE and proximal SAE. This study found no difference in IgM Memory B cell production between distal vs proximal SAE but did find after SAE there is a larger amount of Ig Memory B cells presents compared to splenectomy (25). In August 2022, the EAST guideline provided recommendations against routine post-splenectomy vaccinations in adult trauma patients who have undergone angioembolization for splenic injury because splenic immune function is retained, and infectious complications are not increased (26).

Revaccinations needs have been established by Class II studies of immune antibody levels and efficacy after initial vaccination (3). Patients receiving PCV 20 alone or PCV15 followed by PPSV23 do not require revaccination. If previously vaccinated with PPSV23, revaccination with PCV20 or PCV15 is recommended ≥ 1 year after PPSV23. If previously vaccinated with PCV13, benefits of providing PCV20 or PCV15 have not been evaluated and adults should complete previously recommended PPSV23 series. However, one dose of PCV20 may be used if PPSV23 is not available (27). Based on the current literature, patients who receive the MenACWY polysaccharide diphtheria toxoid conjugate vaccine (Menveo® or MenQuadfi®) should be revaccinated every 5 years. The *Haemophilus influenzae* type b vaccine (Hiberix®) does not require revaccination (12-14).

There is no Class I data identifying the appropriate timing for pre-splenectomy *Haemophilus*, Pneumococcal or Meningococcal vaccination for patients with nonfunctional or diseased spleens. Vaccination two weeks prior to surgery is commonly practiced, but this is supported only by Class III data (16,17). Pre-splenectomy vaccination has been demonstrated to induce antibody formation in both adults and children (33). The types of antibody produced and time to antibody formation (generally 1 to 4 weeks) does vary from patient to patient. (33-35). The antibody titer required to prevent either pneumococcal carriage or disease is unknown and has been extrapolated from data obtained from the literature on *Haemophilus influenzae* titers (36). In the elective splenectomy patient, therefore, vaccination as soon as splenic disease is diagnosed appears prudent to allow time for antibody production (13). The CDC has outlined recommendations for both initial vaccination in the pediatric population as well as booster (revaccination) requirements in patients with an anatomically present, but non-functional spleen (44).

The CDC recommends that asplenic travelers contact an international health clinic or the CDC (www.cdc.gov) to obtain information on disease risks within the intended country of travel. Asplenic travelers should be advised of the increased risk for Meningococcal meningitis and recommendation of the A and C vaccine for all asplenic individuals traveling to sub-Saharan Africa, India, and Nepal.

REFERENCES

1. King H, Schumacher HB. Splenic studies: I. Susceptibility to infection after splenectomy in infancy. *Ann Surg* 1952; 136(2):239-242.
2. Shatz DV. Vaccination practices among North American trauma surgeons in splenectomy for trauma. *J Trauma* 2002; 53:950-956.
3. Rutherford EJ, Livengood J, Higginbotham M, et.al. Efficacy and safety of pneumococcal revaccination after splenectomy for trauma. *J Trauma* 1995; 39:448-452.
4. Dickerman JD. Traumatic asplenia in adults. *Arch Surg* 1981; 116:361-363.
5. Davidson RN, Wall RA. Prevention and management of infections in patients without a spleen. *Clin Microbial Infect* 2001; 7:657-660.
6. Waghorn DJ. Overwhelming infection in asplenic patients: current best practice preventative measures are not be followed. *J Clin Pathology* 2001; 54:214-218.
7. Brigden ML, Pattulo AL. Prevention and management of overwhelming post-splenectomy infection – an update. *Crit Care Med* 1999; 27:836-842.
8. Spickett GP, Bulliore J, Wallis J, et.al. Northern region asplenia register – analysis of first two years. *J Clin Pathology* 1999; 52:424-429.
9. Williams DN, Bhavjot K. Post-splenectomy care strategies to decrease the risk of infection. *Postgrad Med* 1996; 100: 195-8,201,205.
10. Sumaraju V, Smith LG, Smith SM. Infectious complications in asplenic hosts. *Infect Dis Clin N America* 2001; 15:551-565.
11. Styrt B. Infections associated with asplenia: risks, mechanisms, and prevention. *Am J Med* 1990; 88:33N-42N.
12. Prevention and control of meningococcal disease: recommendations of the Advisory Committee on Immunization Practices (AICP). *MMWR* 2000; 49(RR-07):1-3.
13. Prevention and management of infections in patients without a spleen. *Clin Micro and Infect* 2001; 12:657-680.
14. Recommendations of the Advisory Committee on Immunization Practices (AICP): Use of vaccines and immune globulins in persons with altered immunocompetence. *MMWR* 1993; 42(RR-04):1-18.
15. General recommendations on immunization: recommendations of the Advisory Committee on Immunization Practices (ACIP) and the American Academy of Family Physicians (AAFP). *MMWR* 2002; 51(RR-2):1-36.
16. Shatz DV, Romero-Steiner S, Elie CM, et.al. Antibody responses in post-splenectomy trauma patients receiving the 23-valent pneumococcal polysaccharide vaccine at 14 versus 28 days postoperatively. *J Trauma* 2002; 53:1037-1042.
17. Shatz DV, Schinsky MF, Pais LB, et.al. Antibody responses in post-splenectomy trauma patients receiving the 23-valent pneumococcal polysaccharide vaccine at 1 versus 7 versus 14 days after splenectomy. *J Trauma* 1998; 44(6):760-766.
18. Cooker KG, Howard JM, Alvarado AR, et al. Splenic embolization after trauma: An opportunity to improve best immunization practices. *Association for Academic Surgery* 2018: 293-297.
19. Olthof DC, van der Vlies CH, and Goslings JC. Evidence-Based Management and Controversies in Blunt Splenic Trauma. *Curr Trauma Rep* 2017; 3(1):32-37.
20. Skattum J, Titze TL, Dormagen JB, et al. Preserved splenic function after angioembolization of high grade injury. *Injury* 2012; 43(1):62-66.
21. Nakae H, Shimazu T, Miyauchi H, et al. Does splenic preservation treatment (embolization, splenorrhaphy, and partial splenectomy) improve immunologic function and long-term prognosis after splenic injury? *J Trauma* 2009; 67(3):557-563.
22. Bessoud B, Duchosal MA, Siegrist CA, et al. Proximal splenic artery embolization for blunt splenic injury: clinical, immunologic, and ultrasound-Doppler follow-up. *J Trauma* 2007; 62(6):1481-1486.
23. Malhotra AK, Carter RF, Lebman DA, et al. Preservation of splenic immunocompetence after splenic artery angioembolization for blunt splenic injury. *J Trauma* 2010; 69(5):1126-1130.
24. Olthof DC, Lammers AJJ, van Leeuwen EMM, et al. Antibody response to a T-cell-independent antigen is preserved after splenic artery embolization for trauma. *Clin Vaccine Immunol* 2014; 21(11):1500-1504.
25. Foley PT, Kavnoudias HK, Cameron PU, et al. Proximal Versus Distal Splenic Artery Embolisation for Blunt Splenic Trauma: What is the Impact on Splenic Immune Function? *Cardiovasc Intervent Radiol* 2015; 38(5):1143-1151.
26. Freeman JJ, Yorkgitis BK, Haines K, et al. Vaccination after spleen embolization: A practice management guideline from the Eastern Association for the Surgery of Trauma. *Injury* 2022; 53(11):3569-3574.
27. Kobayashi M, Farrar JL, Gierke R, et al. Use of 15-Valent Pneumococcal Conjugate Vaccine and 20-Valent Pneumococcal Conjugate Vaccine Among U.S. Adults: Updated Recommendations of the Advisory Committee on Immunization Practices. Centers for Disease Control and Prevention. 2022 January 28; 71(4):109–117.

28. Kobayashi M, Farrar JL, Gierke R, et al. Use of 15-Valent Pneumococcal Conjugate Vaccine Among U.S. Children: Updated Recommendations of the Advisory Committee on Immunization Practices. Centers for Disease Control and Prevention. 2022 September 16; 71(37):1174–1181.
29. Prevnar-20™ [package insert]. Wyeth Pharmaceuticals LLC; 2021.
30. Pneumovax-23® [package insert]. Merck & Co, Inc; 2011: 1-9.
31. Use of serogroup B meningococcal vaccines in persons age ≥10 years at increased risk for serogroup B meningococcal disease: recommendations of the Advisory Committee on Immunization Practices (ACIP) 2015. *MMWR* 2015; 64(22): 608-612.
32. Trumenba™ meningococcal group B vaccine. [package insert]. Wyeth pharmaceuticals. 2014; 1-10.
33. Kealey GP, Dhungel V, Wideroff MJ, et al. Patient education and recall regarding postsplenectomy immunizations. *Journal of Surgical Research* 2013; 580-585.
34. Bexsero® meningococcal group B vaccine. [package insert]. Novartis Vaccines and Diagnostics 2015: 1-11.
35. Landgren O, Bjoerkholm M, Konradsen HB, et.al. A prospective study on antibody response to repeated vaccinations with pneumococcal capsular polysaccharide in splenectomized individuals with special reference to Hodgkin's lymphoma. *J Intern Med* 2004; 255:644-673.
36. Grimfors G, Soederqvist M, Holm G, et.al. A longitudinal study of class and subclass antibody response to pneumococcal vaccination in splenectomized individuals with special reference to Hodgkin's disease. *Eur J Haematol* 1990; 45:101-108.
37. Schreiber MA, Pusateri AE, Veit BC, et.al. Timing of vaccination does not affect antibody response or survival after pneumococcal challenge in splenectomized rats. *J Trauma* 1998; 45(4):692-697.
38. Werner AM, Katner HP, Vogel R, et.al. Delayed vaccination does not improve antibody responses in splenectomized rats experiencing hypovolemic shock. *Am Surg* 2001; 67(9):834-838.
39. Klinge J, Hammersen G, Scharf J, et.al. Overwhelming post-splenectomy infection with vaccine-type *Streptococcus pneumoniae* in a 12-year-old girl despite vaccination and antibiotic prophylaxis. *Infection* 1997; 25:368-371.
40. Wong WY, Overturf GD, Powars DR. Infection caused by *Streptococcus pneumoniae* in children with sickle cell disease: epidemiology, immunologic mechanisms, prophylaxis, and vaccination. *Clin Infect Dis* 1992; 14:1124-1136.
41. Siber GR, Gorham C, Martin P, et.al. Antibody response to pretreatment immunization and post-treatment boosting with bacterial polysaccharide vaccines in patients with Hodgkin's disease. *Ann Intern Med* 1986; 104:467-475.
42. Hosea SW, Burch CG, Brown EJ, et.al. Impaired immune response of splenectomized patients to polyvalent pneumococcal vaccine. *Lancet* 1981; 1:804-807.
43. Cimaz R, Mensi C, D'Angelo E, et.al. Safety and immunogenicity of a conjugate vaccine against *Haemophilus influenzae* type b in splenectomized and nonsplenectomized patients with Cooley anemia. *J Infect Dis* 2001; 183:1819-1821.
44. CDC: Preventing pneumococcal disease among infants and young children. *MMWR* 2000; 49:1-38.
45. Meningococcal (groups A, C, Y, and W-135) polysaccharide diphtheria toxoid conjugate vaccine: Menactra® [package insert]. Sanofi Pasteur. 2005 Dec: 1-10.
46. Recommended Adult Immunization Schedule – United States, 2015. *MMWR* 2015; 64(4): 91-94.
47. Schimmer JAG, van der Steeg AFW, Zuidema, WP. Splenic function after angioembolization for splenic trauma in children and adults: A systematic review. *Injury* 2016; 47(3):525-530.
48. Foley PT, Kavnaudias H, Cameron PU, et al. Proximal versus distal splenic artery embolization for blunt splenic trauma: what is the impact on splenic immune function. *Cardiovasc Intervent Radiol* 2015: 1143-1151.

POST-SPLENECTOMY PATIENT INFORMATION SHEET

Name: _____

Splenectomy (splee-nek-tuh-mee) is the name of the operation that was done to remove your spleen. The spleen is a fist-sized organ located in the upper left side of your abdomen (belly). The spleen helps you fight infections, get rid of old or damaged red blood cells, and store blood for your body. Because of either disease or damage to your spleen, it had to be removed. You can live without a spleen, but you may be at a higher risk for certain types of blood infection. To help you fight these infections in the future, you have been given the following immunizations (shots):

Vaccine	Initial Date Given	Repeat Doses	Next Due Date
20-valent pneumococcal conjugate vaccine (PCV20, Prevnar 20™) or 15-valent pneumococcal (PCV15, Vaxneuvance™)			
Only if vaccinated with PCV15: 23-valent pneumococcal vaccine (Pneumovax® 23)	≥8 weeks after PCV15		
Meningococcal polysaccharide/diphtheria toxoid conjugate vaccine (Menveo® or MenQuadfi®)		Yes, ≥2 months after initial <u>and</u> every 5 years	
Meningococcal serogroup B (Bexsero® or Trumenba®)		Yes, timing and number based on vaccine given	
Haemophilus influenza type B (Hiberix®)			
Influenza		Yes, annually	

It is important that you go and see a doctor IMMEDIATELY if you have any of the following symptoms:

- Fever
- Chills
- Abdominal pain
- Skin rash, swelling, redness, or infection
- Diarrhea
- Achy or weak feeling
- Cough
- Vomiting

These are signs that you may have an infection. Without your spleen, a small or minor infection may become very serious, and your doctor needs to examine you and possibly start antibiotics to help your body fight the infection. Always check with your doctor before any dental or invasive procedures, as you may need to take antibiotics before the procedure.

The effect of the vaccines in preventing infection varies from patient to patient and depends on the strength of your immune system when the vaccines were given. You will need to be re-immunized (have the shots again) approximately every 5 years for the rest of your life. You should make sure that your doctor has a copy of this information sheet so that they can help remind you when it is time to be re-immunized.

If you or your doctor has questions about the above information, please contact your surgeon:

Surgeon's Name: _____

Surgeon's Phone Number: _____