

DISCLAIMER: These guidelines were prepared by the Department of Surgical Education, Orlando Regional Medical Center. They are intended to serve as a general statement regarding appropriate patient care practices based upon the available medical literature and clinical expertise at the time of development. They should not be considered to be accepted protocol or policy, nor are intended to replace clinical judgment or dictate care of individual patients.

INFERIOR VENA CAVA FILTER USE IN PATIENTS AT HIGH RISK FOR PULMONARY EMBOLISM

SUMMARY

Pulmonary embolism (PE) remains a significant cause of morbidity and mortality in the critically ill surgical or trauma patient. PE may occur even in the presence of appropriate deep venous thrombosis (DVT) prophylaxis. Patients at high risk for PE may benefit from placement of an inferior vena cava (IVC) filter. These devices have been shown to be effective in the prevention of PE and may be placed at the patient's bedside under ultrasound guidance. Temporary IVC filters, placed when the patient is at the highest risk for PE and removed when the patient's risk for PE has decreased, may represent the best balance between prophylaxis and device-related morbidity.

RECOMMENDATIONS

- **Level 1**
 - **None**
- **Level 2**
 - **Prophylactic inferior vena cava (IVC) filters should be placed in the following high-risk patients with contraindications to anticoagulation**
 1. **Age > 55 years with isolated long bone fractures**
 2. **Severe head injury with coma**
 3. **Head injury with long bone fracture**
 4. **Spinal cord injury with paraplegia or quadriplegia**
 5. **Multiple (≥ 2) long bone fractures with pelvic fracture**
 6. **Multiple (≥ 4) long bone fractures**
 7. **Penetrating pelvic venous injury**
 - **Temporary IVC filters should be considered when the risk of PE or contraindications to anticoagulation is anticipated to be less than two (2) weeks.**
 - **IVC filters may be safely placed at the patient's bedside under either fluoroscopic or ultrasound guidance.**
- **Level 3**
 - **To prevent early PE, IVC filters, when indicated, should be placed as soon after injury as possible.**

INTRODUCTION

Deep venous thrombosis (DVT) and pulmonary embolism (PE) remain common, challenging, and often-devastating complications in the surgical or trauma patient. The average incidence of DVT in the general trauma population is 42% (range 18-90%) and the reported incidence of PE in patients with spinal cord injury (SCI) is 10% (range 4%-22%). Up to 4% of injury-related deaths in the U.S. are caused by PE-related "sudden death", frequently in patients that would otherwise have recovered from their injuries.

EVIDENCE DEFINITIONS

- **Class I:** Prospective randomized controlled trial.
- **Class II:** Prospective clinical study or retrospective analysis of reliable data. Includes observational, cohort, prevalence, or case control studies.
- **Class III:** Retrospective study. Includes database or registry reviews, large series of case reports, expert opinion.
- **Technology assessment:** A technology study which does not lend itself to classification in the above-mentioned format. Devices are evaluated in terms of their accuracy, reliability, therapeutic potential, or cost effectiveness.

LEVEL OF RECOMMENDATION DEFINITIONS

- **Level 1:** Convincingly justifiable based on available scientific information alone. Usually based on Class I data or strong Class II evidence if randomized testing is inappropriate. Conversely, low quality or contradictory Class I data may be insufficient to support a Level I recommendation.
- **Level 2:** Reasonably justifiable based on available scientific evidence and strongly supported by expert opinion. Usually supported by Class II data or a preponderance of Class III evidence.
- **Level 3:** Supported by available data, but scientific evidence is lacking. Generally supported by Class III data. Useful for educational purposes and in guiding future clinical research.

PE following development of DVT is one of the most preventable causes of death in hospitalized patients. DVT prophylaxis using either unfractionated / fractionated heparin or intermittent pneumatic compression devices is not 100% protective against DVT formation nor subsequent PE. This is especially true in the critically ill, high-risk patient who may have barriers or contraindications to the use of such methods of prophylaxis such as complex wounds, CNS (brain and spinal cord) or ocular injuries, external fixators, or traction devices. Other considerations include the fact that many patients will require prophylaxis for months to years if not for the remainder of their lifetime, incurring the significant economic impact and potential morbidity of chronic anticoagulation.

The first step in the prevention of PE is the accurate identification of patients at high risk. Various risk factors for PE have been suggested including the following:

- Age > 55 years with isolated long bone fractures
- Severe head injury with coma
- Head injury with long bone fracture
- Spinal cord injury with paraplegia or quadriplegia
- Multiple (≥ 2) long bone fractures with pelvic fracture
- Multiple (≥ 4) long bone fractures
- Penetrating pelvic venous injury

IVC filters have been proven to be a safe and effective treatment to decrease the risk of PE in various patient populations including the critically ill and traumatically injured. Reported complication rates range from 0-24% with patency rates in excess of 90%. Concerns include the safety and long-term effects of these devices, especially in younger patients, for whom the risk of thromboembolism may be time-limited. The recent availability of removable devices may solve some of these problems, offering protection against PE during the early, highest-risk period, while avoiding the potential long-term complications of a permanent filter.

Another issue is when these devices should be placed. Is it appropriate to delay initiation of anticoagulation for 2-3 days to start anticoagulation after splints, casts and external fixators are removed and bleeding risks are less likely? A patient's risk increases within the first several hours after injury with DVT and/or PE frequently being noted within the first 72 hours. Reports exist of PE in the first 24-48 hours.

Cost is another issue. With the ability to place these devices at the patient's bedside under ultrasound guidance, however, the associated costs can be reduced. Other advantages of the bedside approach include avoidance of transporting critically ill patients, avoidance of radiation exposure, and obviating the need for intravenous contrast.

LITERATURE REVIEW

The Eastern Association for the Surgery of Trauma (EAST) has published extensive evidence-based medicine guidelines on the management of DVT in the trauma patient (1). Their evaluation identified that the risk-benefit ratio is favorable in a high-risk trauma patient. No Level I or II recommendations for IVC filter placement in trauma patients were offered. Level III recommendations included placing an IVC filter in a high-risk trauma patient who cannot receive anticoagulation because of increased bleeding risk. High-risk patients were defined as:

- 1) Severe closed head injury (GCS < 8)
- 2) Incomplete spinal cord injury with para- or quadriplegia
- 3) Complex pelvic fractures with associated long-bone fractures
- 4) Multiple long bone fractures.

In summary, although there is no Class I literature to support insertion of an IVC filter in a trauma patient without an established DVT or PE, Class II and III data is starting to accumulate.

Author	Year	Evidence	Findings
Leach (2)	1985	Class II	Review of 201 trauma patients. Negligible morbidity and no mortality. No pulmonary emboli seen.
Rogers (3)	1993	Class III	Retrospective review of 2525 high-risk trauma patients. Four high-risk groups that account for 92% of PE were identified. <ol style="list-style-type: none"> 1. Patients > 55 years of age with isolated long bone fractures 2. Patients with severe head injury and coma 3. Patients with multiple long bone fractures and pelvic fracture 4. Patients with spinal cord injury and paraplegia or quadriplegia Overall incidence of PE was 1%.
Winchell (4)	1994	Class III	Retrospective review of 9721 trauma patients. 0.37% sustained a clinical or autopsy documented PE. Only 22% had a known DVT. 80% of patients with PE were receiving some form of prophylaxis (including 22% who were receiving both pneumatic compression stockings AND subcutaneous heparin). High-risk patient categories included <ol style="list-style-type: none"> 1. Head and spinal cord injury 2. Head and long bone fracture 3. Severe pelvis and long bone fracture 4. Multiple long bone fractures
Rosenthal (5)	1994	Class III	Retrospective case-control study of 151 trauma patients evaluating an aggressive approach to IVC filter placement in high-risk patients. From 1984-1988, 19 of 94 patients (20%) developed DVT despite prophylaxis (mechanical/ subcutaneous heparin). 8 patients developed PE (2 fatal). 15% of patients sustained PE without DVT (3 fatal). No patient sustained PE after filter placement. 23% of patients with ISS>16 developed PE. From 1988-1992, 29 of 67 patients with ISS>16 had filters placed. 13% of all patients developed DVT. Only 1% of patients with ISS>16 developed PE with the more aggressive approach.
Wilson (6)	1994	Class III	Retrospective evaluation of PE in 2525 trauma patients. 6% of patients with traumatic spinal cord injury (SCI) developed PE. Following a more aggressive utilization of IVC filters, no PE has been noted in 15 patients with SCI over a 6-24 month follow-up period.
Khansarinia (7)	1995	Class II	Prospective case-control evaluation of prophylactic IVC filters in 224 patients. 0% incidence of PE in 108 patients with prophylactic IVC filter vs. 6% in 216 historically matched control patients (4% fatal) (p<0.009).
Rodriguez (8)	1996	Class III	Prospective case (40 patients) vs. injury-matched historical control (80 patients) study. PE decreased from 14% to 1% (p=0.02) with prophylactic IVC filter placement. 44% of PE's occurred in the first week.
Gosin (9)	1997	Class III	Prospective case (250 patients) vs. historical control (249 patients) study. Prophylactic IVC filter placement in high-risk trauma patients decreased the PE rate from 4.8% to 1.6% (p=0.045). No clinically evident complications of IVC filter placement were noted.
Rogers (10)	1997	Class II-III	Retrospective review of high-risk orthopedic trauma patients. High-risk injury patterns for PE included: <ol style="list-style-type: none"> 1) Lower extremity fractures (0.62%) 2) Pelvic fractures (1.3%) 3) Pelvic and LE fractures (2.5%) 4) Non-orthopedic trauma patients (0.15%)

			IVC filters were placed in 35 of 940 patients who met 2 or more of the following criteria: 1) Age > 55 years 2) ISS > 16 3) Complex pelvic fractures 4) Long bone and pelvic fractures 5) Lower extremity or pelvic fracture requiring prolonged bedrest Incidence of PE decreased from historical rate of 1% to 0.2% in study population (p<0.04).
Rogers (11)	1998	Class II	Prospective evaluation of IVC filter placement in 792 trauma patients with 35 at high-risk and a contraindication to anticoagulation. No high-risk patient developed PE with a filter in place. 0.25% incidence of PE in trauma patients not deemed to be at high-risk.

Bedside Insertion, Ultrasound Guidance and Temporary Filters

Author	Year	Evidence	Findings
Nunn (12)	1997	Class II	55 patients undergoing bedside IVC filter placement under ultrasound guidance. 89% had successful placement. Failures were mostly due to inability to visualize the right renal vein due to bowel gas. No procedure related mortality and no PE. Four complications (8.2%) included 1 tilted filter, 1 DVT at the needle puncture site, 1 IVC occlusion, and 1 minor filter migration. Estimated annual cost savings were significant (\$69,800-\$118,300). Various other reports confirm the safety, feasibility, and cost effectiveness of this approach (13-16).
Linsenmaier (17)	1998	Class II	Prospective evaluation of 50 temporary IVC filter (Gunther, Gunther Tulip, Antheor) placements. 100% placement success. All temporary filters were removed in 1-12 days (mean 7.3 days). On removal, 18% showed thrombi in the filter. No patients developed a PE with a filter in place. 2 filters migrated and 1 patient developed an IVC thrombosis. 2 filters required femoral venotomy for removal.
Offner (18)	2003	Class II	Prospective evaluation of 44 temporary IVC filter (Gunther Tulip) placements. 84% were in severely injured patients. Filters were in place an average of 14 ± 1 days (range 3-30 days). Three filters could not be retrieved, 2 because of significant clots below the filter and 1 because of abnormal angulation. No complications associated with insertion or retrieval.

Timing of Prophylaxis

Author	Year	Evidence	Findings
Owings (19)	1997	Class III	Retrospective review of 63 trauma patients with PE. 25% of PE's occurred within the first 4 days of injury. 4 patients had their PE (1 fatal) the day following injury.
Carlin (20)	2002	Class III	Retrospective review of 22 trauma patients who developed PE prior to IVC filter placement. On average, PE was diagnosed 4 ± 2 days from admission and 36% occurred in the first 72 hours.

Follow up & Complications

Author	Year	Evidence	Findings
Greenfield (21)	1995	Class III	20-year follow-up of long-term safety and efficacy of IVC filter placement. Data were available for 54% of placements. Mean follow-up was 56.5 months. 93% had a patent insertion site vein. 5% had significant tilting or migration. 2% had a fractured filter strut. No clinical sequelae were noted for tilt, migration or limb fracture. Caval patency was 96%.
Rogers (11)	1998	Class III	Retrospective review of prophylactic IVC filter placement in 132 trauma patients. 3% demonstrated insertion-related thrombosis and 2.3% PE. 36% had follow-up ultrasound examinations. Mean follow-up time 599 days (range 9-1946 days). One asymptomatic IVC thrombosis was detected. 5.5% demonstrated strut malpositioning with a higher incidence of PE in these patients (6.3% vs. 0%; p=0.05).
Langan (22)	1999	Class III	Retrospective review of 160 trauma patients with prophylactic IVC filters. 47% survey response rate and return for examination, duplex ultrasound, and fluoroscopy. Mean follow up was 19.4 months (range 3 to 57 months). The IVC was visualized in 93% and patency was 100% in these patients. Fluoroscopy failed to show any evidence of filter migration. One known clinical PE in 187 patients (0.5%) in whom a filter was inserted.
Sekharan (23)	2000	Class III	Retrospective review of 90 multi-system trauma patients receiving prophylactic IVC filters. 37% returned for evaluation. Mean follow-up was 68 months. 6% demonstrated DVT, 18% lower extremity edema, 0% PE, 0% migration / limb fracture. No IVC thrombosis.
Greenfield (24)	2000	Class III	Retrospective review of IVC filters in 385 trauma patients (249 prophylactic). Long-term outcome was available in 79%. Mean follow up 2.4 years. 2% had insertion site thrombosis and 15.6% DVT. Migration and tilt were rare and clinically and statistically insignificant. IVC patency was 96.5%. 3 PE's (1.5%).
Wojcik (25)	2000	Class III	Retrospective review of 178 trauma patients. 59% returned for follow-up. Mean follow-up 28.9 months. No clinically symptomatic pulmonary emboli. One IVC filter migration (0.95%). One IVC occlusion (0.95%). In the prophylactic group (n=64), 28 (44%) developed a DVT. 11 patients (10.4%) had LE swelling.
Duperier (26)	2003	Class III	Retrospective review of 133 trauma patients receiving IVC filters. 77% had post-insertion duplex studies. 26% had de novo thrombi. No arteriovenous fistulae were noted. No patients developed clinical evidence of IVC occlusion. One patient had a fatal PE.

REFERENCES

1. Rogers FB, Cipolle MD, Velmahos G, Rozycki G. Management of venous thromboembolism in trauma patients. www.east.org.
2. Leach TA, Pastena JA, Swan KG, Tikellis JI, Blackwood JM, Odom JW. Surgical prophylaxis for pulmonary embolism. *Am Surg* 1994; 60:292-295.
3. Rogers FB, Shackford SR, Wilson J, Ricci MA, Morris CS. Prophylactic vena cava filter insertion in severely injured trauma patients: indications and preliminary results. *J Trauma* 1993; 35:637-642.
4. Winchell RJ, Hoyt DB, Walsh JC, Simons RK, Eastman AB. Risk factors associated with pulmonary embolism despite routine prophylaxis: implications for improved protection. *J Trauma*, 1994;37:600-606.
5. Rosenthal D, McKinsey JF, Levy AM, Lamis PA, Clark MD. Use of the Greenfield filter in patients with major trauma. *Cardiovascular Surgery* 1994; 2:52-55.
6. Wilson JT, Rogers FB, Wald SL, Shackford SR, Ricci MA. Prophylactic vena cava filter insertion in patients with traumatic spinal cord injury: preliminary results. *Neurosurgery* 1994; 35 234-239.
7. Khansarinia S, Dennis J, Veldenz H, Butcher JL, Hartland L. Prophylactic Greenfield filter placement in selected high-risk trauma patients. *J Vasc Surg* 1995; 22:231-236.
8. Rodriguez JL, Lopez JM, Proctor MC, Conley JL, Gerndt SJ, et al. Early placement of prophylactic vena caval filters in injured patients at high risk for pulmonary embolism. *J Trauma* 1996; 40:797-804.
9. Gosin JS, Graham AM, Ciocca RG, Hammond JS. Efficacy of prophylactic vena cava filters in high-risk trauma patients. *Ann Vasc Surg* 1997; 11:100-105.
10. Rogers FB, Shackford SR, Ricci MA, Huber BM, Atkins T. Prophylactic vena cava filter insertion in selected high-risk orthopaedic trauma patients. *J Ortho Trauma* 1997; 11:267-72.
11. Rogers FB, Strindberg G, Shackford SR, Osler TM, Morris CS, et al. Five-year follow-up of prophylactic vena cava filters in high-risk trauma patients. *Arch Surg* 1998; 133:406-412.
12. Nunn CR, Neuzil D, Naslund T, Bass JG, Jenkins JM, et al. Cost effective method for bedside insertion of vena caval filters in trauma patients. *J Trauma* 1997; 43:752-58.
13. Tola JC, Holtzman R, Lottenberg L. Bedside placement of inferior vena cava filters in the intensive care unit. *Am Surg*, 1999; 65:833-838.
14. Benjamin ME, Sandager GP, Cohn EJ, Halloran DB, Cahan MA, et al. Duplex ultrasound insertion of inferior vena cava filters in multitrauma patients. *Am J Surg* 1999; 178:92-97.
15. Sato DT, Robinson KD, Gregory RT, Gayle RG, Parent FN, et al. Duplex directed caval filter insertion in multi-trauma and critically ill patients. *Ann Vasc Surg* 1999; 13:365-371.
16. Sing RF, Cicci CK, Smith CH, Messick WJ. Bedside insertion of inferior vena cava filters in the intensive Care unit. *J Trauma* 1999; 47:1104-1107.
17. Linsenmaier U, Rieger J, Schenk F, Rock C, Mangel E, Pfeifer KJ. Indications, management, and complications of temporary inferior vena cava filters. *Cardiovascular and Interventional Radiology*, 1998; 21:464-469.
18. Offner PJ, Hawkes A, Madayag R, Seale F, Maines C. The role of temporary inferior vena cava filters in critically ill surgical patients. *Arch Surg*, 2003; 138:591-595.
19. Owings JT, Kraut E, Battistella F, Cornelius JT, O'Malley R. Timing of the occurrence of pulmonary embolism in trauma patients. *Arch Surg*, 1997; 132:862-867.
20. Carlin AM, Tyburski JG, Wilson RF, Steffes C. Prophylactic and therapeutic inferior vena cava filters to prevent pulmonary emboli in trauma patients. *Arch Surg*, 2002; 137:521-527.
21. Greenfield LJ, Proctor MC. Twenty-year experience with the Greenfield filter. *Cardiovascular Surgery*, 1995; 3:199-205.
22. Langan EM, Miller RS, Casey WJ, Carsten CG, Graham RM et al. Prophylactic inferior vena cava filters in trauma patients at high risk: Follow-up examination and risk/benefit assessment. *J Vasc Surg* 1999; 30:484-490.
23. Sekharan J, Dennis JW, Miranda EF, Hertz JA, Veldenz HC, et al. Long-term follow-up of prophylactic Greenfield filters in multisystem trauma patients. *J Trauma*, 2000; 51:1087-1091.
24. Greenfield LJ, Proctor MC, Michaels AJ, Taheri PF. Prophylactic vena caval filters in trauma: the rest of the story. *J Vasc Surg* 2000; 32:490-497.
25. Wojcik R, Cipolle MD, Fearen I, Jaffe J, Newcomb J et al. Long-term follow-up of trauma patients with a vena caval filter. *J Trauma*, 1999; 49:839-843.
26. Duperier T, Mosenthal A, Swan K, Kaul S. Acute complications associated with Greenfield filter insertion in high-risk trauma patients. *J Trauma* 2003; 54:545-549.