

DISCLAIMER: These guidelines were prepared by the Department of Surgical Education, Orlando Regional Medical Center. They are intended to serve as a general statement regarding appropriate patient care practices based upon the available medical literature and clinical expertise at the time of development. They should not be considered to be accepted protocol or policy, nor are intended to replace clinical judgment or dictate care of individual patients.

## AIR TRAVEL FOLLOWING TRAUMATIC PNEUMOTHORAX

### SUMMARY

The safety of commercial air travel following traumatic pneumothorax has long been a subject of debate with recommendations to patients being based largely upon anecdotal experience. A single Class II study confirms that commercial air travel is safe 14 days following radiographic resolution of a traumatic pneumothorax.

### RECOMMENDATIONS

#### Level 1

- None

#### Level 2

- **Commercial air travel is safe 14 days following radiographic resolution of a traumatic pneumothorax.**
- **A chest radiograph should be obtained immediately prior to air travel to confirm continued resolution of a patient's pneumothorax.**
- **Air travel with a residual pneumothorax may result in respiratory distress in-flight.**

#### Level 3

- None

### INTRODUCTION

Over 1 billion passengers travel by air worldwide each year (1). Limited data exists as to when it is safe for patients who have sustained a traumatic pneumothorax to travel by air. Studies based on military experience with spontaneous pneumothorax in the 1950's and 1960's are widely quoted, but due to the significant pathophysiologic differences between spontaneous and traumatic pneumothorax, this experience is not applicable to patients with traumatic pneumothorax (2,3). The civilian experience with commercial air travel following traumatic pneumothorax is limited to a few anecdotal case reports with a single prospective clinical trial having recently been conducted (4-6).

The concern for allowing such patients to fly stems from the physiology associated with gas containing body cavities during aircraft cabin pressurization. Air trapped within a body cavity (such as a residual pneumothorax) will expand by up to 30% during flight at typical cruising altitudes (7). Changes in barometric pressure have been postulated as being potentially capable of causing a previously sealed pulmonary air leak to reopen, further increasing the patient's risk for recurrent pneumothorax (7). Should a pneumothorax recur during air travel, the flight crew is neither trained nor equipped to handle the respiratory distress that may develop (8,9). Appropriate medical care in such situations is dependent upon the fortuitous presence of a "passenger-physician" and their training and willingness to assist in caring for the passenger in distress. Case reports describing recurrent pneumothorax during air travel and passenger-physician's attempts to treat affected patients have been described (2-5).

As aircraft cabin pressure decreases during ascent, the partial pressure of oxygen also decreases. At a typical cruising altitude of 35,000-45,000 feet, the aircraft cabin is pressurized to the equivalent of 8,000 feet and the arterial oxygen tension of normal passengers will fall to between 55 and 68 mmHg (1). This decrease in arterial oxygenation is usually not noticed by the patient with normal pulmonary function, but can be a major consideration in transport of patients with impaired cardiopulmonary function. Patients with recent pulmonary trauma, such as pneumothorax, rib fractures, or pulmonary contusion, may already have marginal oxygen saturations. With a decrease in arterial oxygen saturation in-flight, oxygenation in such patients may rapidly become marginal or inadequate. Descent to a flight level of less than 12,500 feet, with its attendant increase in cabin pressure and arterial oxygen saturation, has often been

recommended in the event of an in-flight medical emergency, although this is not always possible (1). Diversion to the nearest airport has also been suggested as a possible solution. Although a reasonable option for most transcontinental flights that can usually land within 20-40 minutes, transoceanic flights may be hours from the nearest airport (1). In the event of cardiopulmonary arrest, even 10 minutes may mean the difference between patient survival and death.

Most commercial airlines recognize that they are not equipped to manage such patients should problems arise in-flight and routinely recommend 2-4 weeks as the minimum waiting period before a patient can safely fly. The Air Transport Medicine Committee guidelines are similar suggesting that patients wait a minimum of 2-3 weeks following radiographic resolution of their traumatic pneumothorax before air travel (1). Until recently, these recommendations were based solely upon expert opinion and not clinical trials.

## **LITERATURE REVIEW**

A single Class II study has been performed to address the issue of how soon patients sustaining a traumatic pneumothorax may safely travel by commercial aircraft (6). During a 14-month period, 15 consecutive patients with traumatic pneumothorax were enrolled in a prospective, Institutional Review Board-approved study. Three patients were unwilling to wait 14 days before flying and chose to travel by either car or train. All returned home uneventfully. Of the 12 patients who agreed to wait 14 days before traveling by air, 10 were tourists from Europe whose only mode of transportation was air travel. All had obvious traumatic pneumothoraces by anterior-posterior chest radiograph. All patients had pre-flight chest radiographs performed prior to air travel to confirm continued resolution of their pneumothorax. Ten of the 12 patients met the requirements of the study protocol waiting an average of  $17.5 \pm 4.9$  days following radiographic resolution of their traumatic pneumothorax before flying. All 10 were completely asymptomatic in-flight.

After initially agreeing to the study protocol, two patients subsequently flew prior to completing the 14-day waiting period. The first patient developed a recurrent pneumothorax 3 days into the 14-day waiting period requiring reinsertion of a chest tube. Following re-expansion of his lung, he had a persistent, but stable, apical pneumothorax. He ultimately flew home to Sweden by commercial airline 15 days after chest tube removal and 30 days post-injury with a small pneumothorax. He was accompanied by a physician and was asymptomatic in-flight. Upon arrival in Sweden, he was admitted to a hospital for ongoing rehabilitation. Chest radiograph demonstrated no change in the size of his apical pneumothorax. The second patient waited 14 days following radiographic resolution, but was noted to have a small pneumothorax on her pre-flight chest radiograph. She was not allowed to fly and repeat radiograph 3 days later demonstrated complete resolution of her pneumothorax. She was advised to wait a total of 14 days following pneumothorax resolution, but anxious to return home chose to fly at that time. During her flight, she developed dyspnea, chest pain, nausea, vomiting, and diaphoresis that she likened to the symptoms she had had during her initial traumatic pneumothorax. These symptoms resolved within a few hours of landing in the United Kingdom. She presented to her physician the following day for evaluation, but a chest radiograph was not obtained. She subsequently did well without further sequelae.

Although seemingly a small study, this represents the 14-month experience of a busy Level I trauma center with a large tourist population. The study supports the current recommendations of the Air Transport Medicine Committee suggesting that commercial air travel is safe 14 days after radiographic resolution of a traumatic pneumothorax. Repeat chest radiographs should be obtained immediately prior to air travel to confirm continued resolution of a patient's pneumothorax. Air travel with a residual pneumothorax may result in respiratory distress in-flight.

## REFERENCES

1. Air Transport Medicine Committee, Aerospace Medical Association. Medical Guidelines for Air Travel. *Av Space Environ Med* 1996; 67(10): Suppl, B1-B8.
2. Voge VM, Anthracite R. Spontaneous pneumothorax in the USAF aircrew population: A retrospective study. *Av Space Environ Med* 1986; 57:939-949.
3. Fuchs HS. Incidence of spontaneous pneumothorax in apparently healthy aircrews. *Aerospace Med* 1967; 38:1286-1288.
4. Wallace WA, Wong T, O'Bichere A, Ellis BW. Managing in flight emergencies. *BMJ* 1995; 311:374-376.
5. Dowd KE. Report of death of passenger under treatment by pneumothorax. *JAM* 1945; 16:346-349.
6. Cheatham ML, Safcsak K. Air Travel Following Traumatic Pneumothorax: When Is It Safe? *Am Surg* 1999; 65:1160-1164.
7. AMA Commission on Emergency Medical Services. Medical aspects of transportation aboard commercial aircraft. *JAMA* 1982; 247(7):1007-1011.
8. Rodenberg H. Medical emergencies aboard commercial aircraft. *Ann Emerg Med* 198; 16:1373-1377.
9. Jagoda A, Pietrzak M. Medical emergencies in commercial air travel. *Emerg Med Clin NA* 1997; 15(1):251-260.